

The estimated median

**DIAGNOSTIC
DELAY IS
~10 YEARS**

in adults with HPP*¹



Rethink HPP diagnosis by earlier recognition of hypophosphatasia signs and symptoms and the impact of persistently low ALP adjusted for age and sex ²

*In adults (n=52) median age of earliest reported manifestations of HPP was 37.6 years (min: 0.2, max: 75.2), which preceded median age at diagnosis (47.5 years; min: 0.2, max: 75.2).¹ Data from the Global HPP Registry, sponsored by Alexion Pharmaceuticals, an observational, longitudinal, multinational, long-term study collecting data on HPP diagnosis, history, clinical course, symptoms (including multisystemic aspects of disease) and burden of illness from patients who have a diagnosis of HPP.^{1,3}

ALP, alkaline phosphatase; HPP, hypophosphatasia.

1. Högler W, et al. *BMC Musculoskelet Disord*. 2019;20(1):80. 2. Khan AA, et al. *Osteoporos Int*. 2024;35(3):431–438. 3. Seefried L, et al. *J Bone Miner Res*. 2020;35(11):2171–2178.

The Alexion logo is a registered trademark of Alexion Pharmaceuticals, Inc.

This material is intended for healthcare professionals only.

© 2025 Alexion Pharmaceuticals, Inc. All rights reserved.

GL/UNB-H/0019 | February 2025

ALEXION[®]
AstraZeneca Rare Disease